

## LAMPIRAN

## anemia \* perdarahan Crosstabulation

			perdarahan		Total
			perdarahan	non_ perdarahan	
anemia	anemia	Count	23	17	40
		Expected Count	20.0	20.0	40.0
		% within anemia	57.5%	42.5%	100.0%
		% within perdarahan	60.5%	44.7%	52.6%
		% of Total	30.3%	22.4%	52.6%
non_anemia	non_anemia	Count	15	21	36
		Expected Count	18.0	18.0	36.0
		% within anemia	41.7%	58.3%	100.0%
		% within perdarahan	39.5%	55.3%	47.4%
		% of Total	19.7%	27.6%	47.4%
Total	Total	Count	38	38	76
		Expected Count	38.0	38.0	76.0
		% within anemia	50.0%	50.0%	100.0%
		% within perdarahan	100.0%	100.0%	100.0%
		% of Total	50.0%	50.0%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.900 <sup>b</sup>	1	.168		
Continuity Correction <sup>a</sup>	1.319	1	.251		
Likelihood Ratio	1.908	1	.167		
Fisher's Exact Test				.251	.125
Linear-by-Linear Association	1.875	1	.171		
N of Valid Cases	76				

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 18.00.

## Symmetric Measures

		Value	Asymp. Std. Error <sup>a</sup>	Approx. T <sup>b</sup>	Approx. Sig.
Interval by Interval	Pearson's R	.158	.113	1.377	.173 <sup>c</sup>
Ordinal by Ordinal	Spearman Correlation	.158	.113	1.377	.173 <sup>c</sup>
N of Valid Cases		76			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.

## paritas \* perdarahan Crosstabulation

		perdarahan		Total	
		perdarahan	non_ perdarahan		
paritas	Nulipara	Count	20	9	29
		Expected Count	14.5	14.5	29.0
		% within paritas	69.0%	31.0%	100.0%
		% within perdarahan	52.6%	23.7%	38.2%
		% of Total	26.3%	11.8%	38.2%
primipara		Count	9	9	18
		Expected Count	9.0	9.0	18.0
		% within paritas	50.0%	50.0%	100.0%
		% within perdarahan	23.7%	23.7%	23.7%
		% of Total	11.8%	11.8%	23.7%
multipara		Count	9	20	29
		Expected Count	14.5	14.5	29.0
		% within paritas	31.0%	69.0%	100.0%
		% within perdarahan	23.7%	52.6%	38.2%
		% of Total	11.8%	26.3%	38.2%
Total		Count	38	38	76
		Expected Count	38.0	38.0	76.0
		% within paritas	50.0%	50.0%	100.0%
		% within perdarahan	100.0%	100.0%	100.0%
		% of Total	50.0%	50.0%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	8.345 <sup>a</sup>	2	.015
Likelihood Ratio	8.557	2	.014
Linear-by-Linear Association	8.235	1	.004
N of Valid Cases	76		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 9.00.

## Symmetric Measures

	Value	Asymp. Std. Error <sup>a</sup>	Approx. T <sup>b</sup>	Approx. Sig.
Interval by Interval Pearson's R	.331	.107	3.021	.003 <sup>c</sup>
Ordinal by Ordinal Spearman Correlation	.331	.107	3.021	.003 <sup>c</sup>
N of Valid Cases	76			

- a. Not assuming the null hypothesis.
- b. Using the asymptotic standard error assuming the null hypothesis.
- c. Based on normal approximation.

Range\_umur \* perdarahan Crosstabulation

			perdarahan		Total
			perdarahan	non_ perdarahan	
Range_umur	16-20	Count	3	4	7
		Expected Count	3.5	3.5	7.0
		% within Range_umur	42.9%	57.1%	100.0%
		% within perdarahan	7.9%	10.5%	9.2%
		% of Total	3.9%	5.3%	9.2%
21-25	21-25	Count	13	9	22
		Expected Count	11.0	11.0	22.0
		% within Range_umur	59.1%	40.9%	100.0%
		% within perdarahan	34.2%	23.7%	28.9%
		% of Total	17.1%	11.8%	28.9%
26-30	26-30	Count	15	7	22
		Expected Count	11.0	11.0	22.0
		% within Range_umur	68.2%	31.8%	100.0%
		% within perdarahan	39.5%	18.4%	28.9%
		% of Total	19.7%	9.2%	28.9%
31-35	31-35	Count	2	9	11
		Expected Count	5.5	5.5	11.0
		% within Range_umur	18.2%	81.8%	100.0%
		% within perdarahan	5.3%	23.7%	14.5%
		% of Total	2.6%	11.8%	14.5%
35-40	35-40	Count	3	8	11
		Expected Count	5.5	5.5	11.0
		% within Range_umur	27.3%	72.7%	100.0%
		% within perdarahan	7.9%	21.1%	14.5%
		% of Total	3.9%	10.5%	14.5%
> 40	> 40	Count	2	1	3
		Expected Count	1.5	1.5	3.0
		% within Range_umur	66.7%	33.3%	100.0%
		% within perdarahan	5.3%	2.6%	3.9%
		% of Total	2.6%	1.3%	3.9%
Total	Total	Count	38	38	76
		Expected Count	38.0	38.0	76.0
		% within Range_umur	50.0%	50.0%	100.0%
		% within perdarahan	100.0%	100.0%	100.0%
		% of Total	50.0%	50.0%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	10.840 <sup>a</sup>	5	.055
Likelihood Ratio	11.368	5	.045
Linear-by-Linear Association	1.921	1	.166
N of Valid Cases	76		

a. 4 cells (33.3%) have expected count less than 5. The minimum expected count is 1.50.

## Symmetric Measures

	Value	Asymp. Std. Error <sup>a</sup>	Approx. T <sup>b</sup>	Approx. Sig.
Interval by Interval Pearson's R	.160	.115	1.395	.167 <sup>c</sup>
Ordinal by Ordinal Spearman Correlation	.167	.115	1.454	.150 <sup>c</sup>
N of Valid Cases	76			

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis.

c. Based on normal approximation.